

Fruit Rot. Begin when Disease first appears or runners start to run.

Peppers (7), Eggplant (5) – Maximum applications per Season 6 for Peppers and 7 for Eggplants. Intervals between applications 7-10 days. **For Anthracnose and Fruit Rot** begin as fruit begins to form.

Watermelons (5), Cantaloupes (5) – Maximum applications per Season 8. Intervals between applications 7-10 days. Start applications when Disease first appears.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in original container. Keep in an area inaccessible to children and pets. **DISPOSAL: If Empty:** Do not reuse container. Place in trash or offer for recycling if available. **If Partly Filled:** Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, absorbed through skin or inhaled. Causes moderate eye irritation. Wash thoroughly with soap and water after handling. Avoid breathing dust or spray mist. Avoid contact with skin, eyes or clothing. Home gardeners applying this product must wear long-sleeved shirt, long pants and chemical-resistant gloves. The gloves must be washed thoroughly with soap and water before removing. Clothes must be changed immediately after using this EBDS product and must be laundered separately from other laundry items before reuse.

FIRST AID

Have the product container or label with you when calling a Poison Control Center or doctor, or going for treatment. You may also contact: 1-800-222-1222 or 1-800-900-4044 for emergency medical treatment information.

If Swallowed	<ul style="list-style-type: none"> • Call Poison Control Center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the Poison control Center or doctor. • Do not give anything by mouth to an unconscious person.
If Inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a Poison Control Center or doctor for treatment advice.
If On Skin, Or Clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a Poison Control Center or doctor for treatment advice.
If In Eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses if present, after the first 5 minutes, then continue rinsing eye. • Call a Poison Control Center or doctor for treatment advice.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wastewaters.

PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour, spill or store near heat or open flame.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith. Voluntary Purchasing Groups, Inc., warrants this product to conform to the chemical description on the label and for the purposes stated on the label. The limit of any liability incurred shall be the purchase price paid by the User or Buyer.

Hi-Yield®

Maneb Garden Fungicide

- **A Wettable Powder That Controls Many Diseases On Shrubs, Flowers And Vegetables.**
- **1½ Tablespoons Makes 1 Gallon Of Spray For Shrubs, Flowers and Vegetables.**

ACTIVE INGREDIENTS:

Maneb (Manganese ethylenebisdithiocarbamate) 80%
(Manganese equivalent as metallic 16.5%)

INERT INGREDIENTS: 20%

TOTAL: 100%

Keep Out Of Reach of Children

CAUTION

See Back Panel for Additional Precautionary Statements

NET WEIGHT 9 OZS.

VOLUNTARY PURCHASING GROUPS, INC.

P. O. BOX 460
BONHAM, TEXAS 75418
Visit Us At: www.v-p-g.com
EPA Reg. No. 7401-303
EPA Est. No. 7401-TX-1
15M-14-65U2



DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

If this product is used on a crop, no other product containing a different EBDC active ingredient may be used on the same crop during the same Season.

HI-YIELD® MANEB GARDEN FUNGICIDE is effective for controlling a wide variety of Diseases on Shrubs, Flowers and Vegetables. When used as directed, this product can control listed Diseases and thus help Plants grow stronger, healthier and more productive.

Although this product will control established Diseases, best results are obtained when used as a protective spray on a regular basis. Where higher and lower rates are given, use lower rates of application for protective sprays and higher rates of application when Disease conditions are severe.

SHRUBS AND FLOWERS

Spray to cover all Plant surfaces thoroughly. Apply when Plants are leafed out. Use 1½ Tablespoons per gallon of water to make spray mix.

Roses: Black Spot – Begin when first leaves unfold. Repeat at 7-10 day intervals.

Azaleas, Camellias: Petal Blight – Apply to litter and soil beneath Plants, also to flowers of Azaleas. Begin when flower buds swell. Repeat at 3-5 day intervals.

Geraniums, Hydrangeas, Lilies: Botrytis Blight – Begin when new growth starts, before Disease appears or at first sign of Disease. Repeat at 7-10 day intervals.

Carnations, Pansies: Alternaria Blight or Leaf Spot, Anthracnose, Botrytis Blight – Begin when new growth starts. Repeat weekly.

Dogwood: Anthracnose – Begin when buds open. Repeat when bracts fall, 4 weeks later and in late Summer.

HOME GARDEN VEGETABLES

Use 1½ level Tablespoons per gallon of water. Spray to thoroughly cover upper and lower surfaces of foliage. Repeat application as noted for each crop. Number in () indicates the number of days to observe between last application and harvest for crop.

Tomatoes (5 – Maximum applications per Season 4. Intervals between applications 7-10 days. **For Anthracnose, Early And Late Blight, Grey Leaf Spot (Stemphylium), Septoria Leaf Spot.** Begin application when Tomatoes on first clusters are well-formed. Do not use on young plants in greenhouses.

Beans (Dry Only) (30) – Maximum applications per Season 6. Intervals between applications 5-7 days. **For Downy Mildew and Rust.** Begin when plants are small.

Cucumbers (5), Squash (5) – Maximum applications per Season 8. Intervals between applications 7-10 days. **For Anthracnose, Downy Mildew, Pythium**